

PERSPECTIVE

Ecological Restoration in the Light of Ecological History

Stephen T. Jackson^{1*} and Richard J. Hobbs^{2*}

Ecological history plays many roles in ecological restoration, most notably as a tool to identify and characterize appropriate targets for restoration efforts. However, ecological history also reveals deep human imprints on many ecological systems and indicates that secular climate change has kept many targets moving at centennial to millennial time scales. Past and ongoing environmental changes ensure that many historical restoration targets will be unsustainable in the coming decades. Ecological restoration efforts should aim to conserve and restore historical ecosystems where viable, while simultaneously preparing to design or steer emerging novel ecosystems to ensure maintenance of ecological goods and services.

“[Nature] is ever shaping new forms: what is, has never yet been; what has been, comes not again.”—Johann Wolfgang von Goethe, 1783, *On Nature* (1)

Ecological restoration is rooted in ecological history. To facilitate the recovery of degraded or damaged ecosystems, knowledge of the state of the original ecosystem and what happened to it is invaluable. However, systematic monitoring of ecosystems, whether deeply degraded or nearly pristine, rarely spans more than the past few decades. Restoration ecologists are forced to assess ecological history by indirect means, ranging from documentary sources (e.g., written descriptions, historical photographs, maps, and paintings) to paleoecological records from natural archives (e.g., tree-rings, rodent middens, and sediments of lakes, peatlands, oceans, and estuaries). Fortunately, both documentary and natural archives can provide records of environmental variables and ecosystem properties in many parts of the world.

Restoration ecology looks to ecological history as a means of identifying appropriate restoration targets—the state of the ecosystem before disruption—and assessing sources of damage (e.g., fire suppression, acid rain, and cultural eutrophication). Restoration targets in the “New Worlds” of the Americas, Australia, and Oceania are identified as the “natural” states existing at the time of European discovery and conquest, that is, just before disruptions associated with land clearance, agriculture, grazing, and wildfire control. Ecological history plays a straightforward role in these applications in identifying the natural state of the landscape and constituent ecosystems (2–4), including the range of variability in disturbance and other properties (5–7).

Deeper consideration of ecological history is leading to revision of this approach. First, the notion of “natural” is being redefined based on increasing awareness that pre-European native cultures often

exerted substantial influence on ecosystems, from simple hunting/harvesting to fire management and direct vegetation alteration (8–11). The nature, duration, and intensity of these impacts varied widely in space and time (10, 12), but few terrestrial or estuarine ecosystems escaped some effects of human activity. Second, climate has changed in the past 500 years, owing to natural causes and more recently to human activities (13). For many ecosystems, restoration to a historic standard is anachronistic. The environment has drifted, and so too have the targets. Ecosystems of even the recent past may be unsustainable under an early 21st-century climate. Finally, human activities leave ecological legacies that may be difficult or impossible to override in restoration. These legacies include extinctions (moas, mastodons) and industrial activities (brownfields, mine-lands). Moreover, more subtle human imprints are being revealed, including the *terra preta* soils of the

pre-Columbian Amazon (12) and soil-nutrient mosaics dating to 17th- to 19th-century English settlers in Massachusetts and 2nd- to 3rd-century Roman settlers in France (14, 15). For many parts of Europe, Asia, and Africa, undisturbed landscapes are too remote in time to provide restoration targets, which may instead comprise cultural landscapes (16, 17).

Despite these complications, predisturbance restoration targets remain worthy goals in many contexts. A key task for the future will be to determine where this remains viable and, conversely, where alternative targets must be considered. Historical studies will remain valuable in determining ecosystem structure and function before disruption (2–5) and in assessing the nature and timing of ecosystem responses to disruptions (4–9, 18–20). Paleoecology, together with observational, experimental, and modeling studies, can identify factors that prevent spontaneous or assisted ecosystem recovery once the obvious factors have been eliminated or mitigated.

Paleoecological and paleoenvironmental records spanning the last 10,000 to 20,000 years are now available for much of the globe, most densely in glaciated terrain but also in many other regions (21, 22). These records provide important perspectives for restoration, not all of them comforting. First, environmental and ecological changes are normal; perhaps the most natural feature of the world in which we find ourselves is its continual flux. The past 20,000 years witnessed a transition from a glacial to an interglacial world, with numerous climatic excursions throughout. Few major terrestrial ecosystems have existed in situ for more than the past 12,000 years (23, 24), and most are considerably younger, some arising only within the past few centuries (24). Every terrestrial locale has been occupied by a series of ecosystems—often contrasting in structure and function—since the



Fig. 1. The Big Woods (Minnesota) landscape, dominated by mesic forest, was savanna and prairie until about 1300 C.E., when droughts and consequent fine-fuel reduction led to reduction of surface fires, allowing tree invasion and expansion (27). [Photo: S. T. Jackson]

¹Department of Botany and Program in Ecology, University of Wyoming, Laramie, WY 82071, USA. ²School of Plant Biology, University of Western Australia, Crawley, WA 6009, Australia.

*To whom correspondence should be addressed. E-mail: jackson@uwyo.edu (S.T.J.); rhobbs@cyllene.uwa.edu.au (R.J.H.)

last glacial period. In the long run, no inherent natural ecosystem or landscape configuration exists for any region. Second, a multitude of ecological realizations arise and dissolve as the environment changes. Different species assemblages develop, leading to ecosystems with differing structure and function. The late-glacial “no-analog” communities—assemblages of plants, vertebrates, and insects with no modern counterpart—are the most dramatic example, but community assembly and disassembly are characteristic of the entire Quaternary (25). Third, the paleoecological record provides numerous case studies of multiple, alternative “natural” states, owing to historical contingencies affecting species migration, site colonization, and extirpation (26). These cases are not always subtle, involving contrasting ecosystems (forests, grasslands, woodlands, and steppe) (Fig. 1) (26, 27).

These observations have the potential for setting restoration ecology adrift from its moorings in notions of objectively identifiable natural states of ecosystems. If natural states are elusive, if the environment is always changing and ecosystems are always coming and going, and if multiple realizations are normal, then the premises underlying ecological restoration to a historic standard come under question. Does ecological history render ecological restoration “quaint”?

Ecological restoration finds new moorings in emphasizing restoration of ecosystem function, goods, and services. Restoration ecologists increasingly recognize the ongoing and often inevitable development of novel ecosystems, resulting from species invasions, climate change, land-use legacies, and altered biogeochemical cycles (28, 29). Restoration efforts emphasize managing for change, which is accepted as inevitable, and interventions are directed toward ensuring that desirable ecological goods and services, including aesthetic values, are maintained (7, 30).

The paleoecological record gives restoration ecologists permission to accept environmental and ecological change and to intervene in ways that will foster biodiversity and vital ecosystem functions. In many cases, this will lead to ecosystems unlike those of the past (7, 25, 28). Restored ecosystems may have combinations of species that have never co-occurred. Many such ecosystems will be contingent not only on scientific and societal judgments but also on particular combinations of climate events, disturbances, extinctions, and immigrations (26). As artificial or capricious as these ecosystems may seem, they must be embraced insofar as environmental change is inevitable, multiple ecological realizations are natural, and contingencies and legacies are embedded in virtually all natural ecosystems.

Even in the face of inevitable environmental change and ecological novelty, efforts to conserve and restore historical ecosystems should be con-

tinued and even accelerated in the immediate future. This presents a seeming paradox, given the increasingly anachronistic nature of historical targets. However, preventing damage is more cost effective than trying to repair damage. Furthermore, our understanding of historic ecosystems is typically far greater than for most novel or engineered systems. An unstated aim in restoration is to avoid creating bigger problems than those we seek to solve. Short-term targets of known, historic ecosystems may minimize the risk of making things worse. Restoration efforts might aim for mosaics of historic and engineered ecosystems, ensuring that if some ecosystems collapse, other

Paleoecology will play important roles in all of these efforts. Paleoecological and paleoenvironmental studies inform our understanding of existing and historical ecosystems, determining the circumstances under which they arose, gauging the range of environmental variability they have experienced, and identifying environmental thresholds at which they will require different levels of intervention. By integrating the “reverse monitoring” of paleoecology with conventional “forward monitoring” and targeted experiments, we can diagnose the point(s) at which existing ecosystems will be unsustainable.

At the same time, paleoecological studies will continue to reveal past ecosystem realizations and their properties at local, regional, and global scales. Paleoecological insights, together with modeling, experimentation, and observation, will advance our capacity to engineer ecosystems successfully. Obviously, the more time we purchase by slowing the rates of global change in all its dimensions, the more we increase our capacity for successful adaptation. We face serious risk that global change will outpace our scientific capacity to prescribe adaptive strategies, let alone implement them.




Fig. 2. Contrasting ecosystem trajectories from historic through present to future configurations, indicating degree of change from the historic ecosystem (e.g., physical environment and species pool). Trajectories 1 to 3 indicate systems in three different states today: relatively unchanged (1), moderately altered (2), and severely altered (3). Colored bands indicate costs of restoration to the approximate historic state. Dotted lines represent realistic interventions for each trajectory; pursuit of A is more difficult and expensive than B. For trajectory 3, the only viable option is to slow the rate of change and direct the system to maintain or improve its value in terms of ecosystem services (C). Paleoecology can help assess viability of different levels of intervention by identifying historical states and their range of variability, determining how far existing systems have drifted from these historic states, assessing the thresholds between required levels of intervention, and guiding design of novel and sustainable ecosystems capable of providing ecological goods and services.

functioning ecosystems will remain to build on. In the meantime, we can continue to develop an understanding of how novel and engineered ecosystems function, what goods and services they provide, how they respond to various perturbations, and the range of environmental circumstances in which they are sustainable (28, 29).

Clearly, rapid environmental change renders these tasks daunting, and a major challenge for ecologists is to develop effective means of assessing the status of, and prognosis for, ecosystems in varying states of alteration (Fig. 2). Which historic ecosystems provide viable targets? Under what circumstances will combined forces of climate change, invasive species, and other global-change elements require that alternative ecosystems be considered? Can we develop the tools and wisdom to support these decisions?

References and Notes

1. Translation by T. H. Huxley, *Nature* **1**, 9 (1869).
2. E. Gorham, L. Rochefort, *Wetlands Ecol. Manage.* **11**, 109 (2003).
3. D. A. Burney, L. P. Burney, *Front. Ecol. Environ* **5**, 483 (2007).
4. D. A. Willard, T. M. Cronin, *Front. Ecol. Environ* **5**, 491 (2007).
5. T. W. Swetnam, C. D. Allen, J. L. Betancourt, *Ecol. Appl.* **9**, 1189 (1999).
6. G. Motzkin, D. Foster, in *Forests in Time*, D. R. Foster, J. D. Aber, Eds. (Yale Univ. Press, New Haven, CT, 2004).
7. C. I. Millar, N. L. Stephenson, S. L. Stephens, *Ecol. Appl.* **17**, 2145 (2007).
8. D. M. J. S. Bowman, *New Phytol.* **140**, 385 (1998).
9. J. B. C. Jackson *et al.*, *Science* **293**, 629 (2001).
10. T. R. Vale, Ed., *Fire, Native Peoples, and the Natural Landscape* (Island Press, Washington, DC, 2002).
11. B. A. Black, C. M. Ruffner, M. D. Abrams, *Can. J. For. Res.* **36**, 1266 (2006).
12. M. B. Bush, M. R. Silman, *Front. Ecol. Environ* **5**, 457 (2007).
13. E. Jansen *et al.*, in *Climate Change 2007: The Physical Science Basis*, S. Solomon *et al.*, Eds. (Cambridge Univ. Press, Cambridge, 2007).
14. D. Foster *et al.*, *Bioscience* **53**, 77 (2003).
15. E. Dambrine *et al.*, *Ecology* **88**, 1430 (2007).
16. G. F. Peterken, *Glob. Ecol. Biogeogr.* **8**, 321 (1999).
17. Z. Naveh, *Restor. Ecol.* **13**, 228 (2005).
18. D. E. Schindler, R. A. Knapp, P. R. Leavitt, *Ecosystems* **4**, 308 (2001).
19. C. L. Schelske *et al.*, *Ambio* **34**, 192 (2005).
20. F. M. Chambers, D. Mauquoy, E. W. Cloutman, J. R. G. Daniell, P. S. Jones, *Biodivers. Conserv.* **16**, 2821 (2007).
21. National Oceanic and Atmospheric Administration, National Climatic Data Center, Paleoclimatology, www.ncdc.noaa.gov/paleo/paleo.html.
22. Neotoma Paleoecology Database, www.neotomadb.org.
23. J. W. Williams, B. Shuman, T. Webb III, P. J. Bartlein, P. L. Leduc, *Ecol. Monogr.* **74**, 309 (2004).
24. S. T. Jackson, *J. Veg. Sci.* **17**, 549 (2006).

SPECIAL SECTION

25. J. W. Williams, S. T. Jackson, *Front. Ecol. Environ* **5**, 475 (2007).
26. S. T. Jackson, J. L. Betancourt, R. K. Booth, S. T. Gray, *Proc. Natl. Acad. Sci. U.S.A.* 10.1073/PNAS.0901644106 (2009).
27. B. Shuman, A. K. Henderson, C. Plank, I. Stefanova, S. S. Ziegler, *Ecology* **90**, 2792 (2009).

28. R. J. Hobbs *et al.*, *Glob. Ecol. Biogeogr.* **15**, 1 (2006).
29. T. R. Seastedt, R. J. Hobbs, K. N. Suding, *Front. Ecol. Environ* **6**, 547 (2008).
30. R. J. Hobbs, V. A. Cramer, *Annu. Rev. Environ. Resour.* **33**, 39 (2008).

31. R. H. W. Bradshaw, W. K. Lauenroth, J. A. Lockwood, J. McLachlan, J. T. Overpeck, W. A. Reiners, D. Sax, and J. W. Williams provided discussion and asked difficult questions.

10.1126/science.1172977